Presentation 6 – Jennifer Vasterling

Prospective Assessment of Neurocognition in Future Gulf-Deployed and Gulf-Nondeployed Military Personnel: A Pilot Study

Co-Pls: Jennifer J. Vasterling, Ph.D.
Susan P. Proctor, D.Sc.
Co-I: Robert Kane, Ph.D.
Consultants: Roberta F. White, Ph.D.
Tim Heeren, Ph.D.
MAJ James Ness, Ph.D.

<u>Funding</u>: DoD/VA (USAMRMC-RADIII; VA Medical Research Service)

Scientific Advisory Board

COL Gary Gackstetter, DVM, MPH, Ph.D.

Matthew J. Friedman, MD, Ph.D.

COL Paul Amoroso, MD, MPH


CDR Margaret A.K. Ryan, MD, MPH

Primary Objective

To examine <u>prospectively</u> neurocognitive outcomes related to Iraq deployment in deployed and comparable nondeployed Army troops.

Why Neurocognitive Outcome?

- Concentration and memory problems commonly reported among 1991 Gulf War veterans
- · Adverse impact on daily life
- Some research suggesting CNS dysfunction in 1991 Gulf War veterans.
- Neuropsychological performance as an objective, "portable" screen of CNS integrity.



Why Prospective Assessment?

- Baseline and post-deployment assessments allow documentation of change over time.
- Addresses questions relative to pre-existing conditions.
- Assessment of change helps address interpretation of "subtle" deficits. (That is, what is minor to one person may represent a significant change to another.)

Design

· Prospective, longitudinal

Time 1: Baseline

Time 2: Post-deployment

• 3 primary samples of Army personnel:

Iraq- deployed (n = 600)

Non-deployed (n = 450)

(?) Sinai-deployed (n = 150)

Sample

• <u>Iraq Deployed Sample</u> (n = 600)

4 units:

2 combat/combat support

2 service support

At least 1 unit Guard or Reserve

• Non-deployed Sample (n = 450)

3 units:

1 Active Duty combat/combat support

1 Active Duty service support

1 Guard or Reserve

Variable Domains

Predeployment Variables

Demographics, baseline cognition/mental status, prior trauma exposure, brain & nervous disease/risk factors, perception of unit cohesion, preparedness, and physical health, military variables

Deployment Variables

Deployment status, MOS, unit type, geographic location, objective environmental exposures, combat and stress exposure, self-reported environmental exposures

Neurocognitive Variables

- Functions robust to most acquired brain insults (e.g., vocabulary)
- Functions sensitive to potential deploymentrelated exposures

(e.g., attention, working memory, learning, memory, motor, processing speed)

Data Sources

- Military Health and Personnel Records
- Military environmental exposure and geographic location data
- Self-report (deployment experiences, risk factors, health perception, mood & emotional symptoms)
- Objective neuropsychological performances

Data Analytic Approaches

- Repeated measures multivariate analysis of (co)variance to examine neurocognition over time by deployment group
- Multivariate stepwise regression to identify the unique contributions of independent variables to postdeployment cognitive performance (over deployment)

Units Assessed

- Combat Arms/Combat Support (Active Duty)
 n = 150
 98.7% (of 152)
- Combat Service Support (Active Duty) n = 151 95.6% (of 158)
- Combat Support (National Guard) n = 53 79.1% (of 67)
- Combat Service Support (Active Duty)
 n = 105
 71.4% (of 147)

Participant Characteristics		
Age (yrs)	27.3 (7.2)	18-56
Education (yrs)	12.9 (1.4)	11 - 18
% women	16.6%	
Rank		
% E4 or below	59.0%	
% E5 – E9	35.1%	
% Officers	5.9%	
Race/Ethnicity		
% Caucasian	51.1%	
% African American	24.0%	
% Hispanic	15.0%	
% Other	9.9%	

Summary

- Prospective, longitudinal data collection, including baseline and post-deployment assessments
- Comparison of Iraq-deploying troops to nondeploying troops (and possibly Sinai-deploying troops)
- Primary outcome measure is neurocognitive performance